

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MYTESI safely and effectively. See full prescribing information for MYTESI.

MYTESI™ (crofelemer) delayed-release tablets for oral use

Initial U.S. Approval: 2012

INDICATIONS AND USAGE

MYTESI is an anti-diarrheal indicated for the symptomatic relief of non-infectious diarrhea in adult patients with HIV/AIDS on anti-retroviral therapy. (1)

DOSAGE AND ADMINISTRATION

One 125 mg delayed-release tablet taken orally twice a day, with or without food. (2)

DOSAGE FORMS AND STRENGTHS

Delayed-Release Tablets: 125 mg (3)

CONTRAINDICATIONS

None (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
- DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
 - Risks of Treatments in Patients with Infectious Diarrhea
- ADVERSE REACTIONS
 - Clinical Trials Experience
- DRUG INTERACTIONS
 - Drug Interaction Potential
 - Nelfinavir, Zidovudine, and Lamivudine
- USE IN SPECIFIC POPULATIONS
 - Pregnancy
 - Nursing Mothers
 - Pediatric Use
 - Geriatric Use

WARNINGS AND PRECAUTIONS

Rule out infectious etiologies of diarrhea before starting crofelemer. If infectious etiologies are not considered, there is a risk that patients with infectious etiologies will not receive the appropriate therapy and their disease may worsen. (5.1)

ADVERSE REACTIONS

Most common adverse reactions (incidence \geq 3%) are upper respiratory tract infection, bronchitis, cough, flatulence and increased bilirubin. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Napo Pharmaceuticals at 1-844-722-8256 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- Pregnancy:** Based on animal data, may cause fetal harm. (8.1)
- Pediatric Use:** Safety and effectiveness of MYTESI has not been established in patients less than 18 years of age. (8.4)

See 17 for PATIENT COUNSELING INFORMATION

Revised 06/2016

- Use in Patients with Low CD4 Counts and High Viral Loads
- OVERDOSAGE
- DESCRIPTION
- CLINICAL PHARMACOLOGY
 - Mechanism of Action
 - Pharmacodynamics
 - Pharmacokinetics
- NONCLINICAL TOXICOLOGY
 - Carcinogenesis, Mutagenesis, Impairment of Fertility
- CLINICAL STUDIES
- HOW SUPPLIED/STORAGE AND HANDLING
- PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

INDICATIONS AND USAGE

MYTESI is indicated for symptomatic relief of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy.

DOSAGE AND ADMINISTRATION

The recommended dose of MYTESI is one 125 mg delayed-release tablet taken orally two times a day, with or without food. MYTESI tablets should **not be crushed or chewed**. Tablets should be swallowed whole.

DOSAGE FORMS AND STRENGTHS

MYTESI is a white, oval, enteric-coated 125 mg delayed-release tablet printed on one side with 125SLXP.

CONTRAINDICATIONS

None.

WARNINGS AND PRECAUTIONS

5.1 Risks of Treatment in Patients with Infectious Diarrhea

If infectious etiologies are not considered, and MYTESI is initiated based on a presumptive diagnosis of non-infectious diarrhea, then there is a risk that patients with infectious etiologies will not receive the appropriate treatments, and their disease may worsen. Before starting MYTESI, rule out infectious etiologies of diarrhea. MYTESI is not indicated for the treatment of infectious diarrhea.

ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. A total of 696 HIV-positive patients in three placebo-controlled trials received MYTESI for a mean duration of 78 days. Of the total population across the three trials, 229 patients received a dose of 125 mg twice a day for a mean duration of 141 days, 69 patients received a dose of 250 mg twice a day for a mean duration of 139 days, 102 patients received a dose of 250 mg four times a day for a mean duration of 14 days, 54 patients received a dose of 500 mg twice a day for a mean duration of 146 days, and 242 patients received a dose of 500 mg four times a day for a mean duration of 14 days.

Adverse reactions for MYTESI that occurred in at least 2% of patients and at a higher incidence than placebo are provided in Table 1.

Table 1: Adverse Reactions Occurring in at Least 2% of Patients in the 125 mg Twice Daily Group

Adverse Reaction	Crofelemer 125 mg BID* N = 229 n (%)	Placebo N = 274 n (%)
Upper respiratory tract infection	13 (5.7)	4 (1.5)
Bronchitis	9 (3.9)	0
Cough	8 (3.5)	3 (1.1)
Flatulence	7 (3.1)	3 (1.1)
Increased bilirubin	7 (3.1)	3 (1.1)
Nausea	6 (2.6)	4 (1.5)
Back pain	6 (2.6)	4 (1.5)
Arthralgia	6 (2.6)	0
Urinary tract infection	5 (2.2)	2 (0.7)
Nasopharyngitis	5 (2.2)	2 (0.7)
Musculoskeletal pain	5 (2.2)	1 (0.4)
Hemorrhoids	5 (2.2)	0
Giardiasis	5 (2.2)	0
Anxiety	5 (2.2)	1 (0.4)
Increased alanine aminotransferase	5 (2.2)	3 (1.1)
Abdominal distension	5 (2.2)	1 (0.4)

* Twice daily

Adverse reactions that occurred in between 1% and 2% of patients taking a 250 mg daily dose of MYTESI were abdominal pain, acne, increased aspartate aminotransferase, increased conjugated bilirubin, increased unconjugated blood bilirubin, constipation, depression, dermatitis, dizziness, dry mouth, dyspepsia, gastroenteritis, herpes zoster, nephrolithiasis, pain in extremity, pollakiuria, procedural pain, seasonal allergy, sinusitis and decreased white blood cell count.

Adverse reactions were similar in patients who received doses greater than 250 mg daily.

DRUG INTERACTIONS

7.1 Drug Interaction Potential

In vitro studies have shown that crofelemer has the potential to inhibit cytochrome P450 isoenzyme 3A and transporters MRP2 and OATP1A2 at concentrations expected in the gut. Due to the minimal absorption of crofelemer, it is unlikely to inhibit cytochrome P450 isoenzymes 1A2, 2A6, 2B6, 2C9, 2C19, 2D6, 2E1 and CYP3A4 systemically [see *Clinical Pharmacology* (12.3)].

Nelfinavir, Zidovudine, and Lamivudine

MYTESI administration did not have a clinically relevant interaction with nelfinavir, zidovudine, or lamivudine in a drug-drug interaction trial.

USE IN SPECIFIC POPULATIONS

Pregnancy

Pregnancy Category C

Reproduction studies performed with crofelemer in rats at oral doses up to 177 times the recommended daily human dose of 4.2 mg/kg revealed no evidence of impaired fertility or harm to the fetus. In pregnant rabbits, crofelemer at an oral dose of about 96 times the recommended daily human dose of 4.2 mg/kg, caused abortions and resorptions of fetuses. However, it is not clear whether these effects are related to the maternal toxicity observed. A pre- and postnatal development study performed with crofelemer in rats at oral doses of up to 177 times the recommended daily human dose of 4.2 mg/kg revealed no evidence of adverse pre- and postnatal effects in offspring. There are, however, no adequate, well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

It is not known whether crofelemer is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for adverse reactions in nursing infants from MYTESI, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The safety and effectiveness of MYTESI have not been established in pediatric patients less than 18 years of age.

Geriatric Use

Clinical studies with crofelemer did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently than younger patients.

Use in Patients with Low CD4 Counts and High Viral Loads

No dose modifications are recommended with respect to CD4 cell count and HIV viral load, based on the findings in subgroups of patients defined by CD4 cell count and HIV viral load.

The safety profile of crofelemer was similar in patients with baseline CD4 cell count less than 404 cells/ μ L (lower limit of normal range) (N=388) and patients with baseline CD4 cell counts greater than or equal to 404 cells/ μ L (N=289)


The safety profile of crofelemer was similar in patients with baseline HIV viral loads less than 400 copies/mL (N = 412) and patients with baseline HIV viral loads greater than or equal to 400 copies/mL (N = 278).

OVERDOSAGE

There has been no reported experience with overdosage of crofelemer.

DESCRIPTION

MYTESI (crofelemer) delayed-release tablets is an anti-diarrheal, enteric-coated drug product for oral administration. It contains 125 mg of crofelemer, a botanical drug substance that is derived from the red latex of *Croton lechleri* Müll. Arg. Crofelemer is an oligomeric proanthocyanidin mixture primarily composed of (+)-catechin, (-)-epicatechin, (+)-gallocatechin, and (-)-epigallocatechin monomer units linked in random sequence, as represented below. The average degree of polymerization for the oligomers ranges between 5 and 7.5, as determined by phloroglucinol degradation.



R = H or OH range n = 3 to 5.5

Inactive ingredients: microcrystalline cellulose, croscarmellose sodium, colloidal silicon dioxide, and magnesium stearate.

Coating ingredients: ethylacrylate and methylacrylate copolymer dispersion, talc, triethyl citrate, and white dispersion which contains xanthan gum, titanium dioxide, propyl paraben, and methyl paraben.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Crofelemer is an inhibitor of both the cyclic adenosine monophosphate (cAMP)-stimulated cystic fibrosis transmembrane conductance regulator (CFTR) chloride ion (Cl⁻) channel, and the calcium-activated Cl⁻ channels (CaCC) at the luminal membrane of enterocytes. The CFTR Cl⁻ channel and CaCC regulate Cl⁻ and fluid secretion by intestinal epithelial cells. Crofelemer acts by blocking Cl⁻ secretion and accompanying high volume water loss in diarrhea, normalizing the flow of Cl⁻ and water in the GI tract.

12.2 Pharmacodynamics

Consistent with the mechanism of action of crofelemer (i.e., inhibition of CFTR and CaCC in the GI lumen), data suggest stool chloride concentrations decreased in patients treated with MYTESI (500 mg four times daily) (n=25) for four days relative to placebo (n=24); stool chloride concentrations decreased in both African American patients treated with MYTESI (n=3) relative to placebo (n=5) and non-African American patients treated with MYTESI (n=22) relative to placebo (n=19).

At a dose 10 times the maximum recommended dose, crofelemer does not prolong the QTc interval to any clinically relevant extent.

12.3 Pharmacokinetics

Absorption

The absorption of crofelemer is minimal following oral dosing in healthy adults and HIV-positive patients and concentrations of crofelemer in plasma are below the level of quantitation (50 ng/mL). Therefore, standard pharmacokinetic parameters such as area under the curve, maximum concentration, and half-life cannot be estimated.

Distribution

The distribution of crofelemer has not been determined.

Metabolism

No metabolites of crofelemer have been identified in healthy subjects or patients in clinical trials.

Elimination

The elimination route has not been identified in humans.

Food Effect

Administration of crofelemer with a high-fat meal was not associated with an increase in systemic exposure of crofelemer in healthy volunteers. In the clinical trial, a single 500 mg dose of crofelemer was administered one-half hour before the morning and evening meals. Therefore, crofelemer may be administered with or without a meal.

Drug-Drug Interactions

Results of a crossover study in healthy volunteers showed crofelemer 500 mg administered four times daily for five days had no effect on the exposure of zidovudine and nelfinavir when administered as a single dose. A 20% decrease in lamivudine exposure was also observed in the same study but was not considered to be clinically important.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals have not been performed to evaluate the carcinogenic potential of crofelemer.

Mutagenesis

Crofelemer was negative in the bacterial reverse mutation assay, chromosomal aberration assay, and rat bone marrow micronucleus assay.

Impairment of Fertility

Crofelemer, at oral doses of up to 738 mg/kg/day (177 times the recommended human daily dose of 4.2 mg/kg), had no effects on fertility or reproductive performance of male and female rats.

14 CLINICAL STUDIES

The efficacy of MYTESI 125 mg delayed-release tablets twice daily was evaluated in a randomized, double-blind, placebo-controlled (one month) and placebo-free (five month), multi-center study. The study enrolled 374 HIV-positive patients on stable anti-retroviral therapy (ART) with a history of diarrhea for one month or more. Diarrhea was defined as either persistently loose stools despite regular use of anti-diarrheal medication (ADM) (e.g., loperamide, diphenoxylate, and bismuth subsalicylate) or one or more watery bowel movements per day without regular ADM use.

Patients were excluded if they had a positive gastrointestinal (GI) biopsy, GI culture, or stool test for multiple bacteria (Salmonella, Shigella, Campylobacter, Yersinia, Mycobacterium), bacterial toxin (Clostridium difficile), ova and parasites (Giardia, Entamoeba, Isospora, Cyclospora, Cryptosporidium, Microsporidium), or viruses (Cytomegalovirus). Patients were also excluded if they had a history of ulcerative colitis, Crohn's disease, celiac sprue (gluten-enteropathy), chronic pancreatitis, malabsorption, or any other GI disease associated with diarrhea.

The study had a two-stage adaptive design. In both stages, patients received placebo for 10 days (screening period) followed by randomization to crofelemer or placebo for 31 days of treatment (double-blind period). Only patients with 1 or more watery bowel movements per day on at least 5 of the last 7 days in the screening period were randomized to the double-blind period. Each stage enrolled patients separately; the dose for the second stage was selected based on an interim analysis of data from the first stage. In the first stage, patients were randomized 1:1:1 to one of three crofelemer dose regimens (125, 250, or 500 mg twice daily) or placebo. In the second stage, patients were randomized 1:1 to crofelemer 125 mg twice daily or placebo. The efficacy analysis was based on results from the double-blind portion of both stages.

Each study stage also had a five month period (placebo-free period) that followed the double-blind period. Patients treated with crofelemer continued the same dose in the placebo-free period. In the first stage, patients that received placebo were re-randomized 1:1:1 to one of the three crofelemer dose regimens (125, 250, or 500 mg twice daily) in the placebo-free period. In the second stage, patients that received placebo were treated with crofelemer 125 mg twice daily in the placebo-free period.

The median time since diagnosis of HIV was 12 years. The percentage of patients with a CD4 cell count of less than 404 was 39%. The percentage of patients with a HIV viral load greater than or equal to 1000, 400 to 999, and less than 400 HIV copies/mL was 7%, 3%, and 9%, respectively; the remainder had a viral load that was not detectable. The median time since diarrhea started was 4 years. The median number of daily watery bowel movements was 2.5 per day.

Most patients were male (85%). The percentage of patients that were Caucasian was 46%; the percentage of patients that were African-American was 32%. The median age was 45 years with a range of 21 to 68 years.

In the double-blind period of the study, 136 patients received crofelemer 125 mg twice daily, 54 patients received 250 mg twice daily, 47 patients received 500 mg twice daily, and 138 patients received placebo. The percentages of patients that completed the double-blind period were 92%, 100%, 85%, and 94% in the 125 mg, 250 mg, 500 mg, and placebo arms, respectively.

Most patients received concomitant protease inhibitors (PI) during the double-blind period (Table 2). The most frequently used ARTs in each group were tenofovir/emtricitabine, ritonavir, and lopinavir/ritonavir.

Table 2: Concomitant ART Use in the Double-Blind Period

	125 mg BID (N = 136) n (%)	250 mg BID (N = 54) n (%)	500 mg BID (N = 46) n (%)	Placebo BID N = 138 n (%)
Any ART	135 (99)	53 (98)	45 (98)	134 (97)
Any PI	87 (64)	41 (76)	33 (72)	97 (70)
Tenofovir/ Emtricitabine	45 (33)	22 (41)	16 (35)	52 (38)
Ritonavir	46 (34)	18 (33)	15 (33)	49 (36)
Lopinavir/ Ritonavir	30 (22)	21 (39)	15 (33)	40 (29)
Efavirenz/ Tenofovir/ Emtricitabine	30 (22)	7 (13)	7 (15)	21 (15)
Tenofovir disoproxil fumarate	18 (13)	8 (15)	5 (11)	14 (10)
Atazanavir sulfate	19 (14)	3 (6)	6 (13)	22 (16)
Abacavir w/ lamivudine	17 (13)	5 (9)	5 (11)	18 (13)
Darunavir	19 (14)	4 (7)	4 (9)	14 (10)
Raltegravir	16 (12)	4 (7)	5 (11)	11 (8)
Valaciclovir hydrochloride	12 (9)	8 (15)	5 (11)	16 (12)
Fosamprenavir	12 (9)	6 (11)	4 (9)	13 (9)
Zidovudine w/ lamivudine	12 (9)	3 (6)	3 (7)	15 (11)
Lamivudine	7 (5)	6 (11)	4 (9)	6 (4)
Nevirapine	8 (6)	6 (11)	3 (7)	9 (7)
Atazanavir	5 (4)	6 (11)	2 (4)	2 (1)

Abbreviations: ART = antiretroviral therapy; PI = Protease Inhibitor; BID = twice daily.

The primary efficacy endpoint was the proportion of patients with a clinical response, defined as less than or equal to 2 watery bowel movements per week during at least 2 of the 4 weeks of the placebo-controlled phase. Patients who received concomitant ADMs or opiates were counted as clinical non-responders.

A significantly larger proportion of patients in the crofelemer 125 mg twice daily group experienced clinical response compared with patients in the placebo group (17.6% vs. 8.0%, 1-sided p < 0.01).

In the randomized clinical study, examination of duration of diarrhea, baseline number of daily watery bowel movements, use of protease inhibitors, CD4 cell count and age subgroups did not identify differences in the consistency of the crofelemer treatment effect among these subgroups. There were too few female subjects and subjects with an HIV viral load > 400 copies/mL to adequately assess differences in effects in these populations. Among race subgroups, there were no differences in the consistency of the crofelemer treatment effect except for the subgroup of African-Americans; crofelemer was less effective in African-Americans than non-African-Americans.

Although the CD4 cell count and HIV viral load did not appear to change over the one month placebo-controlled period, the clinical significance of this finding is unknown because of the short duration of the placebo-controlled period.

Of the 24 clinical responders to crofelemer (125 mg twice daily), 22 entered the placebo-free period; 16 were responding at the end of month 3, and 14 were responding at the end of month 5.

16 HOW SUPPLIED/STORAGE AND HANDLING

Crofelemer delayed-release tablets, 125 mg, are white, oval enteric-coated tablets printed on one side with 125SLXP. They are available in the following package size:

Bottles of 60: NDC 70564-802-60

Store at 20°C-25°C (68°F-77°F); excursions permitted between 15°C-30°C (59°F-86°F). See USP Controlled Room Temperature.

17 PATIENT COUNSELING INFORMATION

- Instruct patients that MYTESI tablets may be taken with or without food.
- Instruct patients that MYTESI tablets should not be crushed or chewed. Tablets should be swallowed whole.

Rx Only

Manufactured by Patheon, Inc. for



Napo Pharmaceuticals, Inc., San Francisco, CA 94105

Copyright © Napo Pharmaceuticals, Inc.

US Patent Nos. 7,341,744 and 7,323,195.

NP-367-2 06/16

The botanical drug substance of MYTESI is extracted from *Croton lechleri* (the botanical raw material) that is harvested from the wild in South America.